

Left/right limb judgement task performance following total knee replacement

Rosser, Crystal; Punt, Timothy; Ryan, Cormac

DOI:
[10.3233/BMR-171104](https://doi.org/10.3233/BMR-171104)

License:
None: All rights reserved

Document Version
Peer reviewed version

Citation for published version (Harvard):
Rosser, C, Punt, T & Ryan, C 2018, 'Left/right limb judgement task performance following total knee replacement', *Journal of Back and Musculoskeletal Rehabilitation*. <https://doi.org/10.3233/BMR-171104>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:
Checked for eligibility 13/09/2018

Published in Journal of Back and Musculoskeletal Rehabilitation
DOI: 10.3233/BMR-171104

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Abstract

Purpose: Working body schema (WBS) of the limbs may be indirectly assessed using left/right limb judgement (LRLJ) task performance. This study aimed to investigate if: 1) Total Knee Replacement (TKR) patients perform LRLJ tasks using implicit motor imagery 2) patients have a disrupted WBS following a TKR for the replaced knee compared to the contralateral knee 3) LRLJ task performance changes following post-surgical rehabilitation using change in upper limb LRLJ task performance as a control.

Methods: In a convenience sample ($n=18$, age 69 ± 7 yrs, 12F 6M) of TKR patients <1 month post-surgery, WBS was assessed using LRLJ task performance for the upper (pictures of the hand) and lower limb (pictures of the foot) before and after rehabilitation. Accuracy and response time (RT) were analysed using a series of $2\times2\times2$ ANOVAs.

Results: LRLJ task performance for images corresponding with the operated and non-operated side were comparable for accuracy ($p=0.83$) and RT (0.28). Accuracy for hand images was comparable from baseline to post-rehabilitation ($p=0.54$) whereas accuracy for feet images increased significantly ($p=0.03$). Responses for awkward posture images were significantly slower than for more natural postures ($p=0.001$).

Conclusions: LRLJ task performance data reflected biomechanical constraints that were indicative of implicit motor imagery being performed by patients. There was no evidence of a disrupted LRLJ task performance for the replaced knee compared to the contralateral knee. Following post-surgical rehabilitation patients' lower limb LRLJ task performance improved whilst hand LRLJ task performance remained unchanged. These findings are the first to show that working body schema improves with rehabilitation following TKR, this may explain some of the clinical improvements seen. Implicit motor imagery could theoretically be a useful adjunct to current post-TKR rehabilitation.

Key words: Total knee replacement; working body schema; Implicit motor imagery.

Introduction:

The total knee replacement (TKR) has become an increasingly popular surgical procedure for the treatment of knee osteoarthritis when conservative treatments have failed.¹ While patient outcomes following joint replacement are broadly positive approximately 18% of patients report only a poor to fair outcome² and reports of continuing high levels of pain unrelated to mechanical structural dysfunction are not uncommon.³ The central nervous system may present a novel therapeutic target for maximising the recovery of these patients.

The brain maps that integrate the ~~sensory and~~ motor cortices of different bodily regions have been referred to as working body schema (WBS).⁴ An efficient WBS enables accurate planning of a coordinated movement. There is a growing body of evidence that chronic pain sufferers have a distorted WBS corresponding to the body part in pain.⁵⁻⁶ It is postulated this disruption may play a role in maintaining an individual's pain state in those suffering from a range of chronic pain conditions.⁷⁻¹¹

A disruption in the WBS in those with longstanding knee osteoarthritis (OA) pain has been demonstrated.¹² Using a left/right limb judgement (LRLJ) task performance, where images of feet were presented to the patients who must immediately judge whether the image is of a left or a right foot, the accuracy of left/right judgements was shown to be poorer in those with knee OA compared with healthy controls.¹² WBS disruptions have also been reported in people with hand OA.¹³ This altered WBS may contribute to the maintenance of pain in patients with OA. It is possible such a disruption is present in patients following a TKR, as many will have suffered from OA knee pain for some time prior to surgical intervention. This

disrupted WBS may partially explain why some patients report continuing high levels of pain and dysfunction post-surgery.

To date, no studies have investigated the integrity of the WBS of patients with a TKR or how that WBS changes following a course of rehabilitation. No studies have quantified LRLJ task performance in patients with TKR or the strategies used during the performance of such a task. This study aimed to investigate if: 1) TKR patients perform LRLJ tasks using implicit motor imagery 2) patients have a disrupted WBS following a TKR for the replaced knee compared to the contralateral knee 3) LRLJ task performance changes following post-surgical rehabilitation using change in upper limb LRLJ task performance as a control. Throughout, LRLJ task performance was assessed using pictures of the feet for the lower limb and pictures of the hand for the upper limb.

Methods:

Participants:

A convenience sample of individuals who had received a TKR and were referred to physiotherapy for post-operative rehabilitation were recruited into this study between September 2013 and August 2014. Participants were included if they were ≥ 18 years of age, English speaking, had the capacity to provide informed consent, and had undergone TKR surgery within the past month. Participants were excluded if they had a history of a neurological condition, a history of contralateral TKR or invasive knee surgery within the past six months, a history of foot or ankle surgery (on either limb) in the past six months, an infection of the knee, a visual impairment that would impede ability to complete the LRLJ task performance. All participants provided written informed consent prior to participation. Ethical approval for the study was granted by XXXXXXXX University's School of Health and Social Care Governance and Research Ethics committee [086/13] and The NHS NRES committee North East – Newcastle and North Tyneside 2 [13/NE/0244]. The study was conducted in line with the Declaration of Helsinki.

Procedure:

Participants referred for post-operative rehabilitation were invited to participate in the study within one month of their operation. Baseline data were collected prior to receiving any post-operative rehabilitation. Demographic information was initially collected (age, gender, body mass index, hand/foot dominance, side of TKR, number of days' post-surgery, length of time of knee pain prior to surgery, any current pain in the contralateral knee or upper limbs). Limb dominance was assessed by asking participants which foot they would kick a ball with and which hand they write with. Two knee-specific standardised physical function tests were

carried out to assess active range of motion (AROM) of the operated knee in sitting and the ability to do an active straight leg raise (ASLR) in supine.¹⁴⁻¹⁶ Three self-reported questionnaires were completed; knee pain (average pain in the last 24h, 100mm pain Visual Analogue Scale (VAS) with 0 = no pain and 100 = worst pain imaginable)¹⁷ Knee-Injury Osteoarthritis Score Short-Form (KOOS-PS)¹⁸ and the Euro-Qol 5D-5L (EQ-5D-5L).¹⁹ Each participant then completed a LRLJ task for the upper and lower limb. Following a course of rehabilitation (involving exercise in both a one-to-one and group setting focusing upon active range of motion, strength, balance, and flexibility) participants were invited back for a follow up assessment. All the baseline data was collected again at this point.

Left Right Limb Judgement task:

In the LRLJ task images of the upper limb (hands) and lower limb (feet) were presented to the participant on a computer screen. The participants were required to identify whether the image was a left or right image i.e. left/right judgement. The accuracy and response time (RT) of identification were recorded. The left/right judgements used line drawings presented to the participants via a computer based program (E-Prime® 2.0 Psychology Software Tools, Inc.). The drawings were replicated by permission from the study by Parsons et al²⁰ and consisted of 48 images each depicting a foot in varying laterality (left or right), view (big toe, dorsum, sole or heel) or rotation (0, 60, 120, 180, 240, 300°) of the lower limb. Upper limb images were displayed in five different views of each hand (front, back, little finger, thumb and wrist) and twelve different degrees of rotation (0-330° each separated by 30°).

Participants were seated on a chair with a monitor positioned on a table at eye level (60cm distance) and hands positioned palm down on the table with the index finger of each hand resting on the keyboard (left hand on 'V', right hand on 'N'). A practice run of eight images

was performed (four hands, four feet) to familiarise with the task followed by a further 48 hand images and 48 foot images. This was performed once for lower limb images and once for upper limb images. Each image was presented in the centre of the screen followed by a small fixation cross for a random period between 1,000 and 1,500 ms, images remained on the screen until a response was made. Participants were prompted to keep their head, upper limbs and lower limbs motionless during the task apart from the responding finger.

Data analysis:

Accuracy for each trial reflected the correct or incorrect laterality judgment (i.e. left or right) for each image presented. Response time (RT) was the period in milliseconds from the onset of each image to when a response (key press) was made. Response times faster than 500ms and slower than 10000ms were excluded from the RT analysis; this accounted for less than 1% of all responses. Also, only correct responses were entered for the RT calculations. The median RT for each participant in line with the factors of interest (see below) was then entered for statistical analysis. Accuracy and RT data were analysed using a series of analyses of variance (ANOVA). ~~A 2x2x2 ANOVA with repeated measures, the factors being Limb (hand, foot), Time (Time 1, Time 2) and Side (affected, unaffected) was conducted for Accuracy and for RT.~~ To address aim 1 a further 2x2x2 ANOVA with repeated measures ~~was~~ ere conducted to explore biomechanical constraints across accuracy and RT data; here the factors were Limb (hand, foot), Time (Time 1, Time 2) and Awkwardness (natural, awkward). Biomechanical constraints refer to the established finding in hand and foot-based LRLJ task performance, where the time to recognise the laterality of the limb presented is closely associated with the time it takes to actually move the limb from its current position to the position pictured.²¹ Accordingly, regardless of the degree of rotation away from neutral, response times are slower (and accuracy poorer) for images reflecting more awkward limb

positions from a biomechanical perspective.²²⁻²⁵⁴ These awkwardness effects provide confidence that individuals are mentally rotating their own limb and are therefore considered confirmatory of implicit motor imagery. For these analyses, only images where biomechanical constraints have previously been clearly identified were included. This included all views for images of hands (back, palm, thumb, wrist) but only half the views (sole, big toe) for images of feet.²⁴⁻²⁵⁹ Accordingly, neutral images of left hands (fingers pointing upwards) reflect more natural (medial) postures when rotated clockwise and more awkward (lateral) postures when rotated anti-clockwise, with the reverse pattern for images of right hands. For images of feet, the categorisation of images is based on the same principles though the neutral positions are not always 'toes up'. The categorisation of images followed the influential approach taken by Parsons.^{24xx}

To address aims 2 and 3, a

A 2x2x2 ANOVA with repeated measures, the factors being Limb (hand, foot), Time (Time 1, Time 2) and Side (affected, unaffected), was conducted for Accuracy and for RT.

In addition we undertook a secondary analysis of our data to explore the potential relationship between pain and WBS. Specifically, we carried out correlations between: A) the change in pain (averaged between both legs) and the change in LRLJ task accuracy (for both legs) B) the change in pain for the affected leg and the change in LRLJ task accuracy for the affected leg and C) the change in pain for the unaffected leg and the change in LRLJ task accuracy for the unaffected leg.

Results:

Eighteen individual's provided baseline data [age 68.9±7.3yrs (mean±SD); gender 12F, 6M; BMI 30±6kg.m⁻²; duration of knee pain 42months (24-51months) [median (Interquartile

Formatted: Left

Range)]]]. The average length of time post-operation at baseline was 25days (18-29days) [median (Interquartile Range)]. All participants were right hand and right leg dominant. Four participants reported upper limb pain. Fifteen participants provided follow-up data on completion of their post-operative rehabilitation on average 89days (73-97months97days) [median (Interquartile Range)] post-surgery. From pre to post post-surgical ~~rehabilitation~~ ~~rehabilitationthere was a group trend toward, on average, there wass~~ decreased pain in the operated knee, increased pain in the contralateral knee, improved knee function (operated side) and quality-of-life. The participant characteristics are presented in table 1.

Insert table 1 here

Limb x Time x Side

~~The LRLJ task performance data for the hand and the feet at pre and post rehabilitation are shown in table 2. The accuracy ANOVA revealed a significant main effect of Limb [$F(1,14)=22.26$, $p<0.0003$) and a Limb x Time interaction [$F(1,14)=4.54$, $p=0.05$). There were no other significant main effects or interactions. The simple effects of the interaction showed that accuracy for images of hands was comparable across the two time points (Time 1 = 0.87, Time 2 = 0.88, $p=0.54$) whereas accuracy for images of feet increased significantly between the pre and post testing sessions (Time 1 = 0.68, Time 2 = 0.76, $p=0.03$) (Figure 1).~~

Insert figure 1 here

Insert table 2 here

~~Response times for images of hands (mean=1895ms) were faster than for images of feet (mean=2310ms) leading to a significant main effect of Limb [$F(1,14)=14.49, p=0.002$]. The response time ANOVA revealed no other main effects or interactions.~~

Limb x Time x Awkwardness

Comparing performance for natural vs. awkward images suggested participants demonstrated the typical biomechanical constraints that are a hallmark of implicit motor imagery for limb laterality recognition tasks (See table 23). The related ANOVA for RTs revealed an awkwardness effect [$F(1,14)=16.63, p=0.001$] with responses for images showing more awkward postures (mean = 2455ms) being significantly slower than responses for more natural postures (mean = 1986ms). ~~While there~~ There was a significant main effect for Limb ~~remained here~~ [$F(1,14)=20.32, p=0.0005$], but there were no other significant main effects or interactions (Time, $F(1,14)=3.83, p=0.07$; Limb*Time, $F(1,14)=0.43, p=0.52$; Limb*Awkwardness, $F(1,14)=0.54, p=0.48$; Time*Awkwardness, $F(1,14)=0.22, p=0.65$; Limb*Time*Awkwardness, $F(1,14)=0.26, p=0.62$) suggesting ~~that~~ participants consistently used implicit motor imagery to complete the task for images of both hands and feet across both time points (Figure 12). The corresponding ANOVA for accuracy data again showed ~~no~~ a significant main effect for Limb, $F(1,14)=29.79, p=0.0008$; participants were more accurate in responding to images of hands (mean = 0.9) than feet (mean = 0.69). ~~s~~ There were no other significant main effects or interactions ($Time, F(1,14)=0.30, p=0.59$; Awkwardness, $F(1,14)=1.76, p=0.21$; Limb*Time, $F(1,14)=0.03, p=0.87$; Limb*Awkwardness, $F(1,14)=1.02, p=0.33$; Time*Awkwardness = $F(1,14)=0.72, p=0.41$; Limb*Time*Awkwardness, $F(1,14)=0.03, p=0.87$).

Insert figure 12 here

Insert table ~~23~~ here

Limb x Time x Side

The LRLJ task performance data for the hand and the feet at pre and post rehabilitation are shown in table ~~32~~. The accuracy ANOVA revealed a significant main effect of Limb [$F(1,14)=22.26, p<0.0003$] and a Limb x Time interaction [$F(1,14)=4.54, p=0.05$]. There were no other significant main effects or interactions (Time, $F(1,14)=3.34, p=0.09$; Side, $F(1,14)=0.06, p=0.82$; Limb*Side, $F(1,14)=0.04, p=0.85$; Time*Side, $F(1,14)=0.41, p=0.53$; Limb*Time*Side, $F(1,14)=2.06, p=0.17$). The simple effects of the interaction showed that accuracy for images of hands was comparable across the two time points (Time 1 = 0.87, Time 2 = 0.88, $F(1,14)=0.40, p=0.54$) whereas accuracy for images of feet increased significantly between the pre and post testing sessions (Time 1 = 0.68, Time 2 = 0.76, $F(1,14)=5.77, p=0.03$) (Figure ~~24~~).

Insert figure ~~24~~ here

Insert table ~~32~~ here

Response times for images of hands (mean=1895ms) were faster than for images of feet (mean=2310ms) leading to a significant main effect of Limb [$F(1,14)=14.49, p=0.002$]. The response time ANOVA revealed no other main effects or interactions (Time, $F(1,14)=0.52, p=0.48$; Side, $F(1,14)=0.32, p=0.58$; Limb*Time, $F(1,14)=1.63, p=0.22$; Limb*Side, $F(1,14)=1.60, p=0.23$; Time*Side, $F(1,14)=2.70, p=0.12$; Limb*Time*Side, $F(1,14)=0.76, p=0.40$).

Secondary exploratory analysis

There was no evidence of a relationship between: A) the change in pain (averaged between both legs) and the change in LRLJ task accuracy (for both legs) [A] Spearman's $\rho = 0.12$, $p = 0.40$] B) the change in pain for the affected leg and the change in LRLJ task accuracy for the affected leg [Spearman's $\rho = -0.09$, $p = 0.80$] and C) the change in pain for the unaffected leg and the change in LRLJ task accuracy for the unaffected leg [Spearman's $\rho = 0.04$, $p = 0.90$].

Discussion:

This is the first study to quantify LRLJ task performance in TKR patients. LRLJ task performance data reflected biomechanical constraints that were indicative of implicit motor imagery being performed by patients.^{20,25} This finding provides confidence that participants used implicit motor imagery when making a judgement about the images with which they were presented. Response times and accuracy were comparable for images corresponding with the operated side compared to the unaffected side for the lower limb. Thus there was no evidence of a disrupted WBS for the limb where the knee was replaced compared to the contralateral knee. Finally, over the course of post-surgical rehabilitation, accuracy of judging the laterality of foot images improved from pre to post treatment while the accuracy for hands was unchanged. This suggests that improvements in WBS after lower limb rehabilitation were somatotopically specific (i.e., limited to the lower limb).~~This suggests WBS for the lower limb improves over a period of rehabilitation whilst hand WBS remains unchanged.~~

Performance of the LRLJ task was better for natural rather than awkward postures. Thus, patients demonstrated performance reflecting biomechanical constraints that have become the hallmark of implicit motor imagery when performing LRLJ tasks. It has been demonstrated that patients do not always use this strategy when completing LRLJ task performance.²⁵

When non-implicit motor imagery based strategies are used awkward positions are recognised in a similarly accurate and fast manner to natural positions. Using non-implicit motor imagery strategies theoretically provides little insight into the patients WBS and it would likely be ineffective for enhancing WBS if the LRLJ task performance was being used for motor learning purposes. Our data demonstrates that TKR patients' do use implicit motor imagery strategies when performing LRLJ tasks.

No effect was found for images corresponding with the operated vs. contralateral side for either accuracy or RT. While some studies have shown poorer performance for an impaired or painful side for either response time^{4,8,26} or accuracy^{12,27} reflecting a difficulty with mental rotation of the affected limb, others have reported a more general ~~decline-reduction~~ rather than it being specific to one side.^{12,28-30} In all these cases, LRLJ task performance have been interpreted as patients having difficulty with implicit motor imagery consistent with the findings of the present study. If the LRLJ task performance is measuring the efficiency of the WBS should we expect to see a difference between the affected and unaffected side? The conflicting results in the literature suggest further understanding is required as to why some patient groups demonstrate slower response times compared to controls and others do not and why some studies have demonstrated asymmetric response times between the affected vs. unaffected side and others have shown no asymmetry. The presence of bilateral pain may have been a complicating factor when analysing LRLJ task performance in the present study with respect to side differences. The contralateral knee was nearly as painful pre-rehabilitation and more painful post-rehabilitation than the affected knee. Given that pain has been shown to be associated with impaired LRLJ task performance^{4,8,26,27} it may be that WBS (i.e, task performance) for both knees was impaired. The absence of an age and gender

matched pain free control group prohibited this possibility being explored, which is a limitation of this study.

A change over time in the accuracy scores of the LRLJ task performance was only apparent in the feet ($p < 0.05$). This improvement appears consistent with self-reported function and reflects the improved ability to simulate and actually move the knee/limb post rehabilitation. Hand accuracy scores did not demonstrate a significant change over time suggesting the changes found for feet were indicative of an improvement in the WBS of the knee rather than practice effects of the test itself. Improvement in WBS has been shown to mirror improvements in pain and function in a number of clinical conditions such as phantom limb pain⁷ and complex regional pain syndrome.⁶ The reasons for improvement can only be speculated upon though it is likely that the large amount of physical movement of the knee associated with rehabilitation, and or the reduction in pain, resulted in improved WBS and thus LRLJ task performance. However, we tentatively investigated the potential role of pain reduction in a series of exploratory correlational analysis and found no evidence of a relationship between change in pain and change in LRLJ performance.

~~Performance of the LRLJ task was better for natural rather than awkward postures. Thus, patients demonstrated performance reflecting biomechanical constraints that have become the hallmark of implicit motor imagery when performing LRLJ tasks. It has been demonstrated that patients do not always use this strategy when completing LRLJ task performance.²⁵ When non implicit motor imagery based strategies are used awkward positions are recognised in a similarly accurate and fast manner to natural positions. Using non implicit motor imagery strategies theoretically provides little insight into the patients WBS and it would likely be ineffective for enhancing WBS if the LRLJ task performance was being used~~

~~for motor learning purposes. Our data demonstrates that TKR patients' do use implicit motor imagery strategies when performing LRLJ tasks.~~

Limitations section

The images used to investigate LRLJ knee performance used pictures of the feet. The use of foot images in the LRLJ task performance to reflect the WBS of the knee/whole limb has been adopted in previous studies.¹² It is proposed the mental rotation of the foot will also involve the associated mental rotation of knee. Key strengths of this study was the use of standardised images²⁰ and software (E-Prime 2.0) that allowed millisecond precision. Additionally, the experimenter was present for all testing, ensuring that participants all responded in the same way and maintained the same position throughout testing. Variation in these aspects can modulate data from LRLJ task performance considerably and are a limitation of studies using LRLJ task performance via online data collection. Furthermore, this study identified better LRLJ task performance for hands than feet for both accuracy and response time which is in line with previous literature,³¹⁻³² providing confidence in our data. As this is an observational study no claims of cause and effect can be made. Additionally, the inclusion of a control group of non-TKR patients would have been beneficial.

Clinical implications

Implicit motor imagery could potentially be used in clinical rehabilitation to improve recovery. The underlying mechanisms of implicit motor imagery is such that it activates similar areas in the brain to those activated during actual movement³³, and by doing so without the person experiencing pain, implicit motor imagery aims to un-pair the typically strong temporal association between movement and pain. ~~The underlying mechanisms of implicit motor imagery is such that it activates similar areas in the brain to those activated~~

~~during actual movement³³ and therefore can potentially bypass the output of pain as the threat may be considered lower than if the actual movement was taking place.~~ It has been tentatively suggested that implicit motor imagery could be used as an intervention for TKR patients who have significant pain post-surgery.³⁴ Our data suggest that the WBS may be impaired post-surgery and it has the capacity to improve with standard rehabilitation. Our data also suggest that TKR patients can utilise implicit motor imagery strategies when undertaking LRLJ task performance. Further research is required to fully investigate the presence and extent of any WBS deficit in TKR patients, the potential clinical implications associated with WBS deficit in this patient group and the potential clinical utility of implicit motor imagery as an adjunct to care.

Conclusions

In conclusion, this study provides the first evidence to suggest that WBS of the lower limb ~~may be impaired~~ in people after TKR ~~and~~ selectively improves following rehabilitation (i.e., no improvement in LRLJ task performance for upper limb images). Importantly, the LRLJ lower limb task is valid in this population - demonstrating typical RT and accuracy hallmarks of implicit motor imagery performance. Given the small sample and observational nature of the study, no firm clinical recommendations can be made. Further studies should evaluate whether implicit motor imagery training via LRLJ tasks may be a useful adjunct to current post-TKR rehabilitation given the positive effect of this training in other chronic pain conditions.

References:

1. Gauchard GC, Vancon G, Meyer P, Mainard D, Perrin PP. On the role of knee joint in balance control and postural strategies: Effects of total knee replacement in elderly subjects with knee osteoarthritis. *Gait Posture*. 2010; 32: 155-160.
2. Dakin, H., Gray, A., Fitzpatrick, R., MacLennan, G., Murray, D. Rationing of total knee replacement: a cost-effectiveness analysis on a large trial data set. *BMJ*. 2012; 2: 1-10.
3. Gonzalez, M. H. and Mekhail, A.O. (2007). The failed total knee arthroplasty: evaluation and etiology. *J Am Acad Orthop Surg*. 2007;12: 436-446.
4. Moseley GL. Why do people with complex regional pain syndrome take longer to recognize their affected hand? *Neurology*. 2004; 62: 2182-2186.
5. Pleger B, Tegenthoff M, Ragert P, et al Sensorimotor Returning in Complex Regional Pain Syndrome Parallels Pain Reduction. *Ann Neurol*. 2005; 57: 425-429.
6. Pleger B, Ragert P, Schwenkreis P, et al. Patterns of cortical reorganization parallel impaired tactile discrimination and pain intensity in complex regional pain syndrome. *Neuroimage*. 2006; 32: 503-510.
7. Flor H, Denke C, Schaefer M, Grusser S. Effect of sensory discrimination training on cortical reorganisation and phantom limb pain. *Lancet*. 2001; 357: 1763-1764.
8. Schwoebel J, Friedman R, Duda N, Coslett, HB. Pain and the body schema. *Brain*. 2001; 124: 2098-2104.
9. Schwoebel J, Coslett HB, Bradt J, Friedman R., Dileo C. Pain and the body schema: effects of pain severity on mental representations of movement. *Neurology*. 2002; 59: 775-777.
10. Moseley GL, Zalucki NM, Wiech K. Tactile discrimination, but not tactile stimulation alone, reduces chronic limb pain. *Pain*. 2008; 137: 600-608.
11. Moseley GL, Flor H. Targeting cortical representations in the treatment of chronic pain a review. *Neurorehabil Neural Repair*. 2012; 26: 646-652.

12. Stanton TR, Lin CW, Smeets RJ, Taylor D, Law R, Moseley GL. Spatially defined disruption of motor imagery performance in people with osteoarthritis. *Rheumatology* (Oxford). 2012; 51: 1455-1464.
13. Gilpin HR, Moseley GL, Stanton TR, Newport R. Evidence for distorted mental representation of the hand in osteoarthritis. *Rheumatology*. 2015; 54: 678-682.
14. Rothstein, J.M., Miller, P.J., Roettger, R.F. Goniometric reliability in a clinical setting. *J Am Phys Ther Assoc*. 1983; 10: 1611-1615.
15. Oldmeadow, L.B., McBurney, H., Robertson, V.J. Hospital stay and discharge outcomes after knee arthroplasty: Implications for physiotherapy practice. *Aust J Physiother*. 2002; 48: 117-121.
16. Lavernia, C., D'Apuzzo, M., Rossi, M.D., Lee, D. Accuracy of knee range of motion and assessment after total knee arthroplasty. *J Arthroplasty*. 2008; 23: 85-91.
17. Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain. *Arthritis Care and Res*. 2011; 63: 240-252.
18. Perruccio AV, Lohmander SL, Canizares M, et al. The development of a short measure of physical function for knee OA_KOOS-Physical Function Shortform (KOOS-PS) - an OARSI/OMERACT initiative. *Osteoarthritis Cartilage*. 2008; 16: 542-50.
19. Baker PN, Deehan DJ, Lees D, Jameson S, Avery PJ, Gregg PJ. The effect of surgical factors on early patient-reported outcome measures (PROM) following total knee replacement. *J Bone Joint Surg Br*. 2012; 94: 1058-1066.
20. Parsons LM, Gabrieli JD, Phelps EA, Gazzaniga MS. Cerebrally lateralized mental representations of hand shape and movement. *J Neurosci*. 1998; 18: 6539-6548.
21. Parsons LM. Temporal and kinematic properties of motor behavior reflected in mentally simulated action. *J Exp Psychol Hum Percept and Perform*. 1994; 20: 709-730.

22. Cooper LA, Shepard RN. Mental transformations in the identification of left and right hands. *J Exp Psychol Hum Percept Perform*. 1975; 104: 48-56.
23. Sekiyama K. Kinesthetic aspects of mental representations in the identification of left and right hands. *Percept Psychophys*. 1982; 32: 89-95.
24. Parsons LM. Imagined spatial transformations of one's hands and feet. *Cogn Psychol*. 1987a; 19: 178-241.
25. King R, Johnson MI, Ryan CG, Robinson V, Martin DJ, Punt TD. My foot? Motor imagery-evoked pain, alternative strategies and implications for laterality recognition tasks. *Pain Med*. 2015; 16: 555-557.
26. Coslett HB, Medina J, Klot D, Burkey AR. Mental motor imagery indexes pain: the hand laterality task. *Eur J Pain*. 2010; 14: 1007-1013.
27. Schmid AB, Coppieters MW. Left/right judgment of body parts is selectively impaired in patients with unilateral carpal tunnel syndrome. *Clin J Pain*. 2012; 28: 615-622.
28. Reinersmann A, Haarmeyer GS, Blankenburg M, et al. Left is where the L is right. Significantly delayed reaction time in limb laterality recognition in both CRPS and phantom limb pain patients. *Neurosci Lett*. 2010; 486: 240-245.
29. Nico D, Daprati E, Rigal F, Parsons L, Sirigu A. Left and right hand recognition in upper limb amputees. *Brain*. 2004; 127: 120-132.
30. Fiorio M, Tinazzi M, Aglioti SM. Selective impairment of hand mental rotation in patients with focal hand dystonia. *Brain*. 2006; 129: 47-54.
31. Ionta S, Fourkas AD, Fiorio M, Aglioti SM. Influence of hands posture on mental rotation of hands and feet. *Exp Brain Res*. 2007; 195: 207-217.
32. Parsons LM. Imagine spatial transformation of one's own body. *J Exp Psychol Gen*. 1987; 116: 172-191.

33. Parsons LM. Integrating cognitive psychology, neurology and neuroimaging. *Acta Psychologica*, 2001; 107: 155-181.
34. Toms AD, Mandalia V, Haigh R, Hopwood B. The management of patients with painful total knee replacement. *Bone Joint J*. 2009; 91: 143-150.